

AD-A093 976

MASSACHUSETTS INST OF TECH CAMBRIDGE DEPT OF CHEMISTRY F/G 20/5
PHOTOGENERATION OF VERY ACTIVE HOMOGENEOUS CATALYSTS USING LASE--ETC(U)
DEC 80 J C MITCHENER, M S WRIGHTON N00014-75-C-0880

UNCLASSIFIED

ONR-TR-24

NL

1 of 1

5/9/90



END
DATE
FILMED
2-8
DTIC

UNCLASSIFIED

LEVEL II

(12)

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER ONR-TR-24 ✓	2. GOVT ACCESSION NO. AD-A093976	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Photogeneration of Very Active Homogeneous Catalysts Using Laser Light Excitation of Iron Carbonyl Precursors		5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report
7. AUTHOR(s) James C. Mitchener and Mark S. Wrighton		6. PERFORMING ORG. REPORT NUMBER
8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0880 ✓		9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Massachusetts Institute of Technology Cambridge, MA 02139
10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-051-579		11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia 22217
12. REPORT DATE December 12, 1980		13. NUMBER OF PAGES 21
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; reproduction is permitted for any purpose of the United States Government; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) Distribution of this document is unlimited.		
18. SUPPLEMENTARY NOTES Prepared for publication in the <u>Journal of the American Chemical Society</u> .		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) laser chemistry, photocatalysis, metal carbonyls, hydrosilation, isomerization		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Fe(CO) ₅ and Fe ₃ (CO) ₁₂ can be photoactivated with intense ultraviolet or visible light from a laser to yield catalytic reaction of alkenes. Irradiation for short time periods with intense light allows limits to be placed on the turn-over rates for the photogenerated catalysts. Isomerization of neat 1-pentene, and reaction of neat 1/1 molar ratio HSiEt ₃ /1-pentene mixtures, at 25°C have been studies using Fe(CO) ₅ and Fe ₃ (CO) ₁₂ as the photochemical catalyst precursor. Data show that turnover rates (1-pentene molecules consumed per Fe(CO) ₅ or Fe ₃ (CO) ₁₂ molecule initially present per minute) exceed 1000 in most cases. In		

DTIC
ELECTE

JAN 21 1981

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

(over)

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AD A093976

DDC FILE COPY

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

practical terms, we find that neat 1-pentene solutions can be >25% isomerized to cis- and trans-2-pentene in <5 s with $\sim 10^{-2} \text{M}$ $\text{Fe}(\text{CO})_5$; and 3.66 M/3.66 M, HSiEt_3 /1-pentene can be >25% converted to non- C_5 alkene products in <10s at $<10^{-2} \text{M}$ $\text{Fe}_3(\text{CO})_{12}$. These data establish the photogenerated catalysts from $\text{Fe}(\text{CO})_5$ and $\text{Fe}_3(\text{CO})_{12}$ to be among those having the fastest turnover rates known for homogeneous organometallic catalysts. Conventional xenon flash photolysis experiments (<100 μsec flash) indicate that catalytic action is over within seconds after the light pulse. In such experiments, a single flash can yield up to 50% conversion of neat 1-pentene to products. Intense irradiation does not significantly alter the distribution of products observed compared to previous studies employing low light intensity excitation.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

OFFICE OF NAVAL RESEARCH

13) CONTRACT NO. 0014-75-C-0880

19) TR-37

Task No. NR 051-579

9) TECHNICAL REPORT, NO. 24

1-1 JNK

6) PHOTOGENERATION OF VERY ACTIVE HOMOGENEOUS CATALYSTS
USING LASER LIGHT EXCITATION OF IRON CARBONYL PRECURSORS.

by

10) James C. Mitchener ~~and~~ Mark S. Wrighton

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

11) 29

12) 22
Prepared for publication in the Journal of the American Chemical Society

December 29, 1980

Reproduction in whole or in part is permitted for any
purpose of the United States Government

This document has been approved for public release and
sale; its distribution is unlimited.

Accession for	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<input type="checkbox"/>
By _____	
Distribution/	
Availability Codes	
Dist	Available for Special
A	

Photogeneration of Very Active Homogeneous Catalysts Using Laser Light
Excitation of Iron Carbonyl Precursors

Abstract: $\text{Fe}(\text{CO})_5$ and $\text{Fe}_3(\text{CO})_{12}$ can be photoactivated with intense ultraviolet or visible light from a laser to yield catalytic reaction of alkenes. Irradiation for short time periods with intense light allows limits to be placed on the turnover rates for the photogenerated catalysts. Isomerization of neat 1-pentene, and reaction of neat 1/1 mole ratio HSiEt_3 /1-pentene mixtures, at 25° C have been studied using $\text{Fe}(\text{CO})_5$ and $\text{Fe}_3(\text{CO})_{12}$ as the photochemical catalyst precursor. Data show that turnover rates (1-pentene molecules consumed per $\text{Fe}(\text{CO})_5$ or $\text{Fe}_3(\text{CO})_{12}$ molecule initially present per minute) exceed 1000 in most cases. In practical terms, we find that neat 1-pentene solutions can be >25% isomerized to cis- and trans-2-pentene in <5 s with $\sim 10^{-2} \text{ M}$ $\text{Fe}(\text{CO})_5$; and 3.66 M/3.66M, HSiEt_3 /1-pentene can be >25% converted to non- C_5 alkene products in <10s at $<10^{-2} \text{ M}$ $\text{Fe}_3(\text{CO})_{12}$. These data establish the photogenerated catalysts from $\text{Fe}(\text{CO})_5$ and $\text{Fe}_3(\text{CO})_{12}$ to be among those having the fastest turnover rates known for homogeneous organometallic catalysts. Conventional Xenon flash photolysis experiments (<100 μsec flash) indicate that catalytic action is over within seconds after the light pulse. In such experiments, a single flash can yield up to 50% conversion of neat 1-pentene to products. Intense irradiation does not significantly alter the distribution of products observed compared to previous studies employing low light intensity excitation.

what is the turnover rate for the photogenerated catalysts? Answering such a question is important in establishing whether photoinitiation of catalysis can provide products at a useful rate. But the answer is not straightforwardly obtained, since the actual catalyst in this system is not well-characterized. We have taken an approach that allows us to put some lower limits on the activity of the catalyst.

The observed quantum yield for consumption of 1-pentene according to (1) or (2) significantly exceeds unity¹⁻³ and indicates that an active catalyst is generated photochemically. But the quantum yield at a single light intensity only gives the efficiency for utilizing incident quanta. The quantum yields from $\text{Fe}(\text{CO})_5$ or $\text{Fe}_3(\text{CO})_{12}$ photocatalyzed reactions are not found to be infinite; catalysis does not persist in the dark after photoinitiation in these systems. Sealed reaction vessels lead to the possibility of the catalyst back reacting with photoejected CO, while purging with an inert gas to remove the CO leads to clusters that are catalytically inactive. This chemistry complicates the determination of catalyst turnover rate. Thus, the approach we have taken has been to determine the observed rate of 1-pentene consumption using very high light intensity excitation where the catalyst turnover rate, not photogeneration of the catalyst, could become the rate limiting feature. The results are given below.

Photocatalysis experiments according to equations (1) and (2) were carried out using an Ar ion laser as the excitation source to provide very high light intensity. Consumption of 1-pentene as a function of irradiation time proves to be a very fast process. Typically, we have monitored the chemistry by gas chromatography, but in these experiments we found that the reaction is so fast that we became concerned about the

finite time interval involved in opening hermetically sealed ampules to quench any residual dark catalytic activity.

Irradiation of several samples for equal time followed by analysis of each at a different time after irradiation reveals no dark after effects; that is, it would appear that when the light is turned off catalysis ceases essentially instantly. The time resolution of this sort of experiment is <30 s, the time needed to quench any catalytic activity by exposing the solutions to O_2 by opening the hermetically sealed ampules. We have found that neat 1-pentene solutions can be quantitatively analyzed by near-infrared absorption spectroscopy. Figure 1 shows the results of catalysis on a portion of the near infrared spectrum of 1-pentene. Independent measurement of the spectra of pure cis-2-pentene and pure trans-2-pentene in the same wavelength range shows that the spectral changes shown in Figure 1 are consistent with the chemistry represented by equation (1) where the ratio of trans-2-pentene to cis-2-pentene is roughly 3 to 1. The near-infrared spectrum of 1-pentene shows bands not present in the 2-pentenenes and allows quantitative analysis of 1-pentene disappearance on a time scale much faster than gas chromatography. Comparison of a near infrared spectrophotometric analysis and gas chromatographic analysis of 1-pentene disappearance show that the near infrared is a reliable quantitative technique. Monitoring the unique near infrared absorbance of 1-pentene shows that catalysis persists for <5 s after irradiation of $Fe(CO)_5$ /1-pentene mixtures is terminated.

Flash photolysis followed by monitoring at 1330 nm shows that catalysis is over within a few seconds after the light is switched off (vide infra).

Table I summarizes typical data for photocatalysis experiments using high light intensity continuous laser excitation. The shortest convenient irradiation time was 5 s in such experiments and roughly 25-50% conversion of neat starting material to product could be obtained using either $Fe(CO)_5$

or $\text{Fe}_3(\text{CO})_{12}$. In reaction of HSiEt_3 with 1-pentene the 2-pentenes account for only ~25% of the 1-pentene reacted; that is, 75% of the 1-pentene yields n-pentane and Si-containing material. In all cases the observed quantum yield, Φ , (number of 1-pentene molecules consumed per incident photon) exceeded unity and turnover rates generally exceeded 1000. The turnover rates given are based on the concentration of $\text{Fe}(\text{CO})_5$ or $\text{Fe}_3(\text{CO})_{12}$ initially present and are thus lower limits. The data shown establish the catalyst from $\text{Fe}_3(\text{CO})_{12}$ or $\text{Fe}(\text{CO})_5$ to be among the most active homogeneous organometallic catalysts.

Determining turnover rates accurately depends on knowing the catalyst concentration and being able to simultaneously measure the reaction rate. Flash photolysis provides a way to generate a certain catalyst concentration and catalytic chemistry could be monitored in the near infrared. We find that catalytic chemistry is easily observable using flash photolysis, and we had hoped that we could monitor the decline of the 1330 nm near infrared overtone absorption following the excitation flash of <100 μs duration. Quantitative analysis, Table I, shows that a single flash can yield a large extent conversion. The difficulty is that flash photolysis of $\text{Fe}(\text{CO})_5$ or $\text{Fe}_3(\text{CO})_{12}$ in 1-pentene yields a near infrared absorbing transient with a lifetime of about one second precluding a quantitative measure of the 1330 nm absorbance. The conclusion to be drawn now is that the catalytic chemistry is over within a few seconds after the excitation flash.⁶ Continued effort is being directed to the use of flash photolysis to measure turnover rates.

Intense irradiation, and especially pulsed excitation, raises the possibility of unusually active species being formed.⁶ For example, the primary result of irradiating $\text{Fe}(\text{CO})_5$ is believed to be that represented by equation (3).⁷ When pulsed excitation is used the leading edge of the



pulse may effect the process represented by equation (3) while the trailing portion may effect further CO loss as in equation (4). The Fe(CO)_4



is known to be photosensitive with respect to CO loss in low temperature matrices.⁸ Photolysis of Fe(CO)_5 in the gas phase can even yield Fe atoms.^{9,10} The consequences of such multiple coordinative unsaturation are not known but we note no new products nor any change in the distribution of products in the photocatalyzed reactions of alkenes represented by equations (1) and (2). We have used a pulsed Nd-Yag laser (frequency tripled to yield 355 nm pulses) excitation of Fe(CO)_5 with <6 nsec pulses, 20 mJ/pulse, and still find no unusual effects on catalytic product distributions. Likewise, 532 nm (<7 nsec, 40 mJ/pulse) irradiation from a frequency doubled Nd-Yag laser does not alter products from $\text{Fe}_3(\text{CO})_{12}$ photocatalysis.

To summarize our main findings, we conclude that photolysis of Fe(CO)_5 or $\text{Fe}_3(\text{CO})_{12}$ yields an exceedingly active catalyst. The active catalyst is likely not an excited state species since the excited lifetimes of Fe(CO)_5 or $\text{Fe}_3(\text{CO})_{12}$ are too short to effect catalysis.⁹ The active species is likely an unsaturated ground state species photoproduced by loss of CO from Fe(CO)_5 or Fe-Fe bond cleavage in $\text{Fe}_3(\text{CO})_{12}$.¹⁰ Very high turnover rates for alkene isomerization and hydrosilation have been found. Moreover, very high turnover numbers (moles of product per mole of Fe initially present) are achieved. For example, when a single flash yields ~50% conversion of neat 1-pentene using $\sim 10^{-2}$ M Fe(CO)_5 the turnover number exceeds 400. Subsequent flashes yield an even greater extent conversion to ultimately completely equilibrate the linear pentenes.⁹ High light intensity does not appear to alter the product distribution where a number of products are possible (alkene + HSiEt_3). The data show that very high light intensity can be used to produce product at a high observed rate and with high quantum efficiency ($>>1$).

Acknowledgements. We thank the Office of Naval Research for partial support of this research. Use of the M.I.T. Spectroscopy Laboratory laser facilities is gratefully acknowledged. M.S.W. acknowledges support as a Dreyfus Teacher-Scholar grant recipient 1975-1980.

James C. Mitchener and Mark S. Wrighton*

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

References

1. Schroeder, M. A.; Wrighton, M.S. J. Am. Chem. Soc., 1976, 98, 551 and J. Organometal. Chem., 1977, 128, 345.
2. (a) Asinger, F.Y.; Fell, B.; Schrage, K. Chem. Ber., 1965, 98, 372, 381;
(b) Asinger, F.; Fell, B.; Collin, G. ibid., 1963, 96, 716;
(c) Carr, M.D.; Kane, V.V.; Whiting, M.C. Proc. Chem. Soc., London, 1964, 408.
3. Austin, R. G.; Paonessa, R. S.; Giordano, P.J.; Wrighton, M.S. Adv. Chem. Ser., 1978, 168, 189.
4. (a) Howard, J.F.; Chalk, A.J. "Organic Synthesis via Metal Carbonyls", Vol. 2, Wender, I.; Pino, P., eds., Wiley: New York, 1977, p. 680-694;
(b) Frankel, E.N.; Emken, E.A.; Peters, H.M.; Davison, V.L.; Butterfield, R.O. J. Org. Chem., 1964, 29, 3292 and 3620.
5. (a) James, B. R. "Homogeneous Hydrogenation", Wiley: New York, 1973;
(b) Taqui Khan, M.M.; Martell, A.E. "Homogeneous Catalysis by Metal Complexes", Vol. I and II, Academic Press: New York, 1974 and references therein.
6. For additional information concerning the nature of the catalyst and its lifetime after photogeneration see the accompanying communication, Chase, D. B.; Weigert, F.J. J. Am. Chem. Soc., 1981, 103, 0000.
7. Geoffroy, G.L.; Wrighton, M.S. "Organometallic Photochemistry", Academic Press: New York, 1979.
8. (a) Poliakoff, M. J. Chem. Soc., Dalton, 1974, 210; (b) Poliakoff, M., Turner, J.J. J. Chem. Soc., Faraday, 1974, 70, 93.
9. Callear, A.B.; Oldman, R.J. Nature, 1966, 210, 730.
10. Korny, Z.; Naaman, R.; Zare, R.N. Chem. Phys. Lett., 1978, 59, 33.

Table I. $\text{Fe}(\text{CO})_5$ and $\text{Fe}_3(\text{CO})_{12}$ Photocatalyzed Reactions of 1-Pentene.^a

A. Continuous Laser Irradiation.^b

Catalyst Precursor (conc., M)	Irrdn. λ , nm (Power, W)	Irrdn. Time, s	Reaction	[1-pentene], M ^c	% 1-Pentene ^d consumed	ϕ^e	Turnover Rate 1-pentene/ catalyst-min ^f
$\text{Fe}(\text{CO})_5$ (8×10^{-3})	333,351,364 (2.5)	5	isomerization	9.14 (neat)	27	68	3700
$\text{Fe}_3(\text{CO})_{12}$ (5×10^{-3})	514.5 (1.3)	10	isomerization	9.14 (neat)	35	44	2400
$\text{Fe}(\text{CO})_5$ (1×10^{-2})	333,351,364 (2.5)	30	hydrosilation	3.66	7	11	700
$\text{Fe}_3(\text{CO})_{12}$ (5×10^{-3})	514.5 (7.5)	10	hydrosilation	3.66	24	13	900
		5	hydrosilation	3.66	42	42	1700
		10	hydrosilation	3.66	50	24	1000
		20	hydrosilation	3.66	42	5	1800
					59	3	1300

B. Flash Photolysis Excitation.^g

Catalyst Precursor (conc., M)	Reaction	[1-pentene], M ^c	% 1-pentene ^d consumed
$\text{Fe}(\text{CO})_5$ (1.4×10^{-2})	isomerization	9.14 (neat)	46
$\text{Fe}(\text{CO})_5$ (1.1×10^{-2})	hydrosilation	3.66	23
$\text{Fe}_3(\text{CO})_{12}$ (5.1×10^{-3})	hydrosilation	3.66	25

^aReactions were generally carried out at 25°C; stirred, 1.0 ml of freeze-pump-thaw degassed solution was irradiated in hermetically sealed Pyrex ampules. Analysis for reaction was by gas chromatography or by near infrared spectrophotometry (see text). Product ratios for isomerization were determined using a 30' x 1/8" 20% Propylene Carbonate on Chromosorb P column operated at 20°C and hydrosilation products were quantitated using a 30' x 1/8" β , β' -ODPN column operated at 50°C. Control experiments show that no photoreaction occurs in the absence of $\text{Fe}(\text{CO})_5$ or $\text{Fe}_3(\text{CO})_{12}$; none is expected since 1-pentene and Et_3SiH are transparent to light transmitted by Pyrex.

Table I. (continued)

The irradiation source was either a Spectra Physics Model 164 Ar ion laser or a Coherent Radiation CR18 Ar ion laser tuned to the indicated wavelength and power. The beam diameter was ~2 mm.

For isomerization reactions the 1-pentene was used as solvent. For hydrosilation HSiEt_3 and 1-pentene were used in a 1/1 mole ratio (3.66 M/3.66 M) without any diluent but using n-hexane as an internal standard for gas chromatographic analyses.

Data shown are for total amount of 1-pentene consumed. For isomerization the only products are trans- and cis-2-pentene but in the presence of HSiEt_3 the products also include n-pentane, (n-pentyl) SiEt_3 , and three isomers of (pentenyl) SiEt_3 . The per cent of 1-pentene that reacts to form n-pentane and Si-containing material is ~75% under the conditions used; i.e. isomerization accounts for only ~25% of 1-pentene disappearance.


Number of 1-pentene molecules consumed per photon incident on the sample.

Turnover rate is given assuming all $\text{Fe}(\text{CO})_5$ or $\text{Fe}_3(\text{CO})_{12}$ forms a catalyst during the period indicated. Thus, these are lower limits on the turnover rate.

Flash photolysis was done using a Xenon Corp. Model 710 apparatus and excitation was one 10 kV, 2000 J, <100 μsec flash of a 1.0 ml degassed sample in a Pyrex ampule.

Figure Caption

Near infrared spectral changes accompanying ultraviolet irradiation ($355 \text{ nm} \pm 20 \text{ nm}$, $\sim 2 \times 10^{-6} \text{ ein/min}$) of 2.0 ml of $1.86 \times 10^{-2} \text{ M Fe(CO)}_5$ in neat 1-pentene. Features with an arrow pointing down are associated with the 1-pentene that is consumed; features with an arrow pointing up are associated with the cis- and trans-2-pentene products formed in a $\sim 1:2.8$ ratio.



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Area Office Attn: Dr. George Sandoz 715 Broadway New York, New York 10005	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Western Regional Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Eastern/Central Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GENNo.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. Paul Delahay Department of Chemistry New York University New York, New York 10003	1	Dr. P. J. Hendra Department of Chemistry University of Southampton Southampton SO9 5NH United Kingdom	1
Dr. E. Yeager Department of Chemistry Case Western Reserve University Cleveland, Ohio 41106	1	Dr. Sam Perone Department of Chemistry Purdue University West Lafayette, Indiana 47907	1
Dr. D. N. Bennion Department of Chemical Engineering Brigham Young University Provo, Utah 84602	1	Dr. Royce W. Murray Department of Chemistry University of North Carolina Chapel Hill, North Carolina 27514	1
Dr. R. A. Marcus Department of Chemistry California Institute of Technology Pasadena, California 91125	1	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
Dr. J. J. Auborn Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. C. E. Mueller The Electrochemistry Branch Materials Division, Research & Technology Department Naval Surface Weapons Center White Oak Laboratory Silver Spring, Maryland 20910	1
Dr. Adam Heller Bell Laboratories Murray Hill, New Jersey 07974	1	Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1
Dr. T. Katan Lockheed Missiles & Space Co, Inc. P.O. Box 504 Sunnyvale, California 94088	1	Dr. J. Boechler Electrochimica Corporation Attention: Technical Library 2485 Charleston Road Mountain View, California 94040	1
Dr. Joseph Singer, Code 302-1 NASA-Lewis 21000 Brookpark Road Cleveland, Ohio 44135	1	Dr. P. P. Schmidt Department of Chemistry Oakland University Rochester, Michigan 48063	1
Dr. B. Brummer EIC Incorporated 55 Chapel Street Newton, Massachusetts 02158	1	Dr. F. Richtol Chemistry Department Rensselaer Polytechnic Institute Troy, New York 12181	1
Library P. R. Mallory and Company, Inc. Northwest Industrial Park Burlington, Massachusetts 01803	1		

TECHNICAL REPORT DISTRIBUTION LIST, 359

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. A. B. Ellis Chemistry Department University of Wisconsin Madison, Wisconsin 53706	1	Dr. R. P. Van Duyne Department of Chemistry Northwestern University Evanston, Illinois 60201	1
Dr. M. Wrighton Chemistry Department Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1	Dr. B. Stanley Pons Department of Chemistry University of Alberta Edmonton, Alberta CANADA T6C 2G2	1
Larry E. Plew Naval Weapons Support Center Code 30736, Building 2906 Crane, Indiana 47522	1	Dr. Michael J. Weaver Department of Chemistry Michigan State University East Lansing, Michigan 48824	1
S. Ruby DOE (STOR) 600 E Street Washington, D.C. 20545	1	Dr. R. David Rauh EIC Corporation 55 Chapel Street Newton, Massachusetts 02158	1
Dr. Aaron Wold Brown University Department of Chemistry Providence, Rhode Island 02192	1	Dr. J. David Margerum Research Laboratories Division Hughes Aircraft Company 3011 Malibu Canyon Road Malibu, California 90265	1
Dr. R. C. Chudacek McGraw-Edison Company Edison Battery Division Post Office Box 28 Bloomfield, New Jersey 07003	1	Dr. Martin Fleischmann Department of Chemistry University of Southampton Southampton SO9 5NH England	1
Dr. A. J. Bard University of Texas Department of Chemistry Austin, Texas 78712	1	Dr. Janet Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. M. M. Nicholson Electronics Research Center Rockwell International 3370 Miraloma Avenue Anaheim, California	1	Dr. R. A. Osteryoung Department of Chemistry State University of New York at Buffalo Buffalo, New York 14214	1
Dr. Donald W. Ernst Naval Surface Weapons Center Code R-33 White Oak Laboratory Silver Spring, Maryland 20910	1	Mr. James R. Moden Naval Underwater Systems Center Code 3632 Newport, Rhode Island 02840	1

TECHNICAL REPORT DISTRIBUTION LIST, 359

<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
1	Dr. R. Nowak Naval Research Laboratory Code 6130 Washington, D.C. 20375	1 Dr. John Kincaid Department of the Navy Strategic Systems Project Office Room 901 Washington, DC 20376
1	Dr. John F. Houlihan Shenango Valley Campus Pennsylvania State University Sharon, Pennsylvania 16146	1 M. L. Robertson Manager, Electrochemical Power Sonics Division Naval Weapons Support Center Crane, Indiana 47522
1	Dr. M. G. Sceats Department of Chemistry University of Rochester Rochester, New York 14627	1 Dr. Elton Cairns Energy & Environment Division Lawrence Berkeley Laboratory University of California Berkeley, California 94720
1	Dr. D. F. Shriver Department of Chemistry Northwestern University Evanston, Illinois 60201	1 Dr. Bernard Spielvogel U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709
1	Dr. D. H. Whitmore Department of Materials Science Northwestern University Evanston, Illinois 60201	1 Dr. Denton Elliott Air Force Office of Scientific Research Bldg. 104 Bolling AFB Washington, DC 20332
1	Dr. Alan Bewick Department of Chemistry The University Southampton, SO9 5NH England	1
1	Dr. A. Himy NAVSEA-5433 NC #4 2541 Jefferson Davis Highway Arlington, Virginia 20362	

TECHNICAL REPORT DISTRIBUTION LIST, 051A

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. M. A. El-Sayed Department of Chemistry University of California, Los Angeles Los Angeles, California 90024	1	Dr. M. Rauhut Chemical Research Division American Cyanamid Company Bound Brook, New Jersey 08805	1
Dr. E. R. Bernstein Department of Chemistry Colorado State University Fort Collins, Colorado 80521	1	Dr. J. I. Zink Department of Chemistry University of California, Los Angeles Los Angeles, California 90024	1
Dr. C. A. Heller Naval Weapons Center Code 6059 China Lake, California 93555	1	Dr. D. Haarer IBM San Jose Research Center 5600 Cottle Road San Jose, California 95143	1
Dr. J. R. MacDonald Chemistry Division Naval Research Laboratory Code 6110 Washington, D.C. 20375	1	Dr. John Cooper Code 6130 Naval Research Laboratory Washington, D.C. 20375	1
Dr. G. B. Schuster Chemistry Department University of Illinois Urbana, Illinois 61801	1	Dr. William M. Jackson Department of Chemistry Howard University Washington, DC 20059	1
Dr. A. Adamson Department of Chemistry University of Southern California Los Angeles, California 90007	1	Dr. George E. Walraffen Department of Chemistry Howard University Washington, DC 20059	1
Dr. M. S. Wrighton Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1		